

Death and long-term disability after gun injury: a cohort analysis

Sheharyar Raza MD, Deva Thiruchelvam MSc, Donald A. Redelmeier MD MS(HSR)

Abstract

Background: Gun injury accounts for substantial acute mortality worldwide and many others survive with lingering disabilities. We investigated whether additional health losses beyond mortality can also arise for patients who survive with long-term disability.

Methods: We conducted a population-based individual patient analysis of adults injured by firearms who had received emergency medical care in Ontario, Canada, from Apr. 1, 2002, to Apr. 1, 2019. Longitudinal cohort analyses were evaluated through deterministic linkages of individual electronic patient files. The primary outcome was death or subsequent application for long-term disability in the years after hospital discharge.

Results: In total, 8313 patients were injured from firearms, of which 3020 were injured from intentional incidents and 5293 were injured from unintentional incidents. A total of 2657 (88.0%) patients with intentional gun injury and 5089 (96.1%) patients with unintentional gun injury survived initial injuries. After a mean 7.75 years of follow-up, patients surviving intentional injuries had a disability rate twice as high as patients surviving unintentional injuries (19.7% v. 10.1%, $p < 0.001$), equivalent to a hazard ratio of 2.01 (95% confidence interval 1.80–2.25). The higher risk of long-term disability for survivors after intentional gun injury was not explained by demographic characteristics, extended to survivors treated and released from the emergency department, and was observed regardless of whether the incident was self-inflicted or from interpersonal assault. Half of the disability cases were identified after the first year. Additional predictors of long-term disability included a lower socioeconomic status, an urban home location, arrival by ambulance transport, a history of mental illness and a diagnosis of substance use disorder.

Interpretation: Our study shows that gun death statistics underestimate the extent of health losses from long-term disability, particularly for those with intentional injuries. Additional and sustainable follow-up medical care might improve patient outcomes.

Guns can be useful for hunting excursions, sport shooting and self-defence. Canadians living in wilderness regions rely on guns for securing food and protecting themselves against animal attacks.¹ However, a large downside of guns is the risk of injury. In Canada, mortality from gun injuries amounts to 800 total deaths annually, equivalent to a rate of 23 deaths per million per year.² An estimated 700 fewer Canadians would die from gun injury each year if per capita mortality rates in Canada matched those in the United Kingdom.^{3,4} Some countries have a higher mortality from gun injuries, including the United States with a rate of about 200 deaths per million population per year.⁵

Many people survive gun injury, which means mortality rates may underestimate the total health losses.⁶ For example, some patients with through-and-through brain injuries (i.e., from a bullet that has passed through leaving entry and exit wounds) stay alive but reside in long-term care institutions.⁷ The intensity of pain or neurologic deficits associated with gun injury can be severe and lasting.⁸ Up to half of those with gun injury show anxiety, depression or other signs of stress while in hospital.^{9,10} Disfigurement can lead to further complications.¹¹ Conversely, some patients view survivorship as a source of

personal pride, community prestige or divine intervention.^{12,13} However, aside from reports after military combat, rigorous studies are near-silent about long-term prognosis and instead focus on acute care survival.^{14–16}

The aim of this study was to examine whether the risks of death and long-term disability are substantial after gun injury, contrary to the rapid recoveries depicted in some action movies. Moreover, we hypothesized that intentional gun injury, relative to unintentional gun injury, would lead to a greater burden of long-term disability because of the differences in wound anatomy, patient characteristics, injury circumstances, trauma severity, counterfactual reasoning, psychological outrage and community supports.^{17–19} Herein we explore this distinction and apply population-wide health care databases to examine long-term outcomes for patients who survive gun injury.

Competing interests: None declared.

This article has been peer reviewed.

Correspondence to: Donald Redelmeier, dar@ices.on.ca

CMAJ Open 2020. DOI:10.9778/cmajo.20190200

Methods

Setting

Ontario is Canada's most populous province, and had a population of 13 069 182 in 2010 (study midpoint) distributed over 1 074 845 km² of land area (urban and rural).^{20,21} Health services data could be analyzed through individually linked population-based databases at ICES.^{22–27} Emergency care was universally available with no user fees for all 178 hospitals in the province and the data could be tracked through encrypted linked ICES databases.^{24,28,29} Prevailing laws during the study period (Apr. 1, 2002, to Apr. 1, 2019) included a mandate for medical reporting of all patients who experienced gun injury.³⁰ Diverse disability programs were available for adults older than 18 years, based on physician assessments that were included in ICES databases.³¹ The population average incidence of disability is about 7 per 1000 people annually in this region.³²

Gun injury

We identified adults aged 16 years or older injured by firearms and who had received emergency medical care between Apr. 1, 2002, and Apr. 1, 2019. These dates included all the years for which data were available in ICES databases and a minimum 1-year follow-up for nearly every patient. Past studies indicate that ICES databases are comprehensive (covering > 99% of emergency departments), connected (linkage rates > 95%) and consistent (diagnostic reliability > 90% compared with chart abstraction).^{25,33,34} Diagnoses were based on the International Classification of Diseases as validated and used in past studies (codes W32–34, X72–74, X93–95 and Y22–24).^{35–39} We excluded those dead at the scene, those living outside of Ontario, those lacking a valid health card number and youth younger than 16 years (eligibility age for disability support). Patients with more than 1 injury were analyzed by first presentation to avoid statistical artifacts from trauma recidivism.⁴⁰

Additional characteristics

Data on patient age (years), sex (binary), socioeconomic status (quintile) and home location (urban, rural) were based on linked demographic databases.^{26,41,42} Additional databases identified incident time (year, season, hour), firearm type (hand gun, long gun, uncertain) and whether the patient arrived by ambulance (air or ground).²⁹ Intentional injuries (self-inflicted, assault) were defined by diagnostic codes with uncertain cases presumed unintentional.^{43,44} We further searched outpatient databases in the prior year to identify earlier psychiatric illnesses and diagnosis of substance use disorder.⁴⁵ General health care use indicators in the prior year also included total hospital admissions, emergency visits and outpatient contacts as proxies for comorbidity.^{46,47}

Acute care

We examined short-term clinical outcomes for secondary descriptive analyses to corroborate past studies.^{18,48–50} Hospital mortality included patients who died in the emergency department, during initial hospital admission or after transfer

to a specialized trauma centre.⁵¹ Hospital length of stay indicated the total time in days from arrival in the emergency department to death, discharge or departure (including patients who left against medical advice).⁵² The number of operations, use of transfusion products and days in intensive care were identified, taking into account those who had none. The available ICES databases lacked information on race, injury circumstances, bullet calibre, vital signs, imaging scans, functional status, formal education and criminal records.⁵³

Primary outcome

The primary outcome was subsequent death or long-term disability (i.e., submission of a new disability support application) for patients who survived initial injury. We considered disability as the dominant outcome along with death as a competing risk (each component also tested in secondary analysis). Such disability support applications included a medical report (Health Status Report, Activities of Daily Living Index, Special Necessities Benefit Form) and involved the patient's physician. In turn, the submission of a medical report allowed tracking of applications using Ontario Health Insurance Plan codes (K050–K054).^{54,55} The available databases lacked information on social supports and financial programs accessed by patients.

We defined disability as the submission of a new disability support application because the document was available, measurable, authenticated and incorporated the patient's perspective.⁵⁵ These techniques have been validated in past research yet can undercount disability because of patients who are reluctant to complain or hesitate to file a claim.⁵⁶ We defined the date of disability by the submission of the application because this time was available, objective and served as a marker for the realization that full recovery might not occur. The available databases did not contain information on the reasons for disability, supports received or how it directly connected to the original gun injury.

Statistical analysis

In our primary comparison, we tested whether the risk of disability in the years following gun injury is higher for those with intentional injuries compared with those with unintentional injuries. We defined the follow-up interval as starting on the day of hospital departure and included only those who survived initial injuries. We used unadjusted cumulative incidence curves to evaluate survivors for death or disability during the decade following injury.⁵⁷ We further examined the relative risk of death and disability for the full cohort of survivors before and after adjusting for additional measured baseline characteristics using the Fine and Gray model of competing risks (subdistributional hazard ratios used as the estimate of relative risks).^{58–61} No data imputation methods were used.

Ethics approval

The study protocol was approved by the Sunnybrook Research Ethics Board, including a waiver for direct individual patient consent.

Results

In total, 8313 individuals were injured by firearms and received emergency medical care during the 17-year study period. Of the total injuries, 63.7% were unintentional injuries and 36.3% were intentional injuries (Table 1). Both groups mainly consisted of men younger than 30 years, and the individuals were widely distributed across socioeconomic status quintiles. Patients with intentional injuries, relative to those with unintentional injuries, tended to live in a city, have an uncertain weapon type and arrive by ambulance transport. Injuries occurring on the weekends were frequent in both groups and a nighttime incident was disproportionately frequent for those with intentional injuries (Appendix 1A, available at www.cmajopen.ca/content/8/3/E469/suppl/DC1). A minority in both groups had a past hospital admission or a past mental health diagnosis.

Acute care outcomes

A total of 2430 patients were admitted to hospital (Table 2). Among those admitted to hospital, about two-thirds required a surgical procedure, one-third required critical care, and a quarter of those admitted to hospital received transfusion products. In total, the cohort accounted for 6191 days in critical care and 24 577 days of hospital stay.

The general profile of short-term acute hospital care suggested a greater severity of injury for patients with intentional incidents compared with those with unintentional incidents, as measured by hospital admission rates, surgical procedures, critical care, blood transfusions, mean days in hospital and risk of acute death (Table 2). A total of 2657 patients with intentional gun injury and 5089 patients with unintentional gun injury survived initial injuries.

Subsequent risk of disability

The 7746 total survivors accounted for 60 098.6 patient-years of follow-up (mean 7.75 yr). Patients surviving intentional gun injury accounted for 584 subsequent cases of disability over 17 669.4 patient-years of follow-up (mean 6.65 yr), equal to an incidence of 33 per 1000 patients annually. Patients surviving unintentional gun injury accounted for 639 subsequent cases of disability over 42 429.2 patient-years of follow-up (mean 8.34 yr), equal to an incidence of 15 per 1000 patients annually. Taken together, intentional gun injury was associated with an increased risk of subsequent disability relative to unintentional gun injury (disability rate 19.7% v. 10.1%, respectively, $p < 0.001$; hazard ratio [HR] 2.01, 95% confidence interval [CI] 1.80–2.25). For both groups, half the disability cases appeared after the first year (Figure 1).

Additional predictors

The risk of subsequent disability associated with gun injury was also related to patient characteristics. Lower socioeconomic status, an urban home location, a nighttime incident time and ambulance arrival were each associated with higher risks (Table 3). A past diagnosis of mental illness or of substance use disorder or a recent emergency department visit

was also associated with higher risks. Conversely, patient age and sex were not significant predictors. The day of the week or the season were also not significant predictors. Adjustment for all measured patient characteristics suggested that intentional gun injury was associated with an increased risk of subsequent disability relative to unintentional gun injury (HR 1.40, 95% CI 1.24–1.60).

Secondary analyses

The higher relative risk of subsequent disability associated with intentional gun injury relative to those with unintentional gun injury extended to important subgroups. In particular, a higher risk was observed for patients treated and released from the emergency department and for patients admitted to hospital (Appendix 1B). Similarly, a higher risk was observed regardless of a history of mental illness, substance use disorder, firearm weapon type, ambulance involvement, health care in the prior year or length of hospital stay. A higher risk was consistent for incidents that were self-inflicted and for incidents from interpersonal assault. No subgroup showed contrary findings and all subgroups with more than 1000 patients showed a significant higher relative risk.

Subsequent mortality

The higher risks of subsequent disability associated with intentional gun injury were also observed for long-term mortality, although the absolute counts were modest (Figure 2). Patients surviving intentional gun injury accounted for 137 subsequent deaths, equal to an incidence of 8 per 1000 patients annually. Patients surviving unintentional gun injury accounted for 156 subsequent deaths, equal to an incidence of 4 per 1000 patients annually. Taken together, intentional gun injury was associated with an increased risk of mortality relative to unintentional gun injury (HR 1.85, 95% CI 1.47–2.33). Additional predictors of subsequent mortality included older age, lower socioeconomic status, an incident occurring at night and a diagnosis of mental illness (Table 3).

Interpretation

We studied gun injury to assess the long-term risks of disability. We found that most patients survived during the initial hospital stay and many later became disabled. The risk of long-term disability was substantial, amounting to more than 1 in 5 patients with intentional injuries and 1 in 10 patients with unintentional injuries. The higher relative risk of long-term disability following intentional gun injury was not fully explained by baseline patient demographic characteristics, occurred regardless of whether the incident was self-inflicted or from interpersonal assault, and extended to those who were not admitted to hospital. Taken together, these data suggest that mortality statistics underestimate the effects of gun injury because many patients do not lose their lives, but lose their livelihoods.^{62,63}

Our study supports past reports of long-term outcomes after other forms of injury. A retrospective cohort study in a Baltimore population evaluated all adult trauma patients over

Table 1: Characteristics of 8313 patients with intentional or unintentional gun injury who received emergency medical care during the study period

Characteristic	No. (%) of patients	
	Intentional* n = 3020	Unintentional n = 5293
Demographic		
Age, yr		
< 30	1857 (61.5)	3178 (60.0)
≥ 30	1163 (38.5)	2115 (40.0)
Sex		
Male	2754 (91.2)	4756 (89.9)
Female	266 (8.8)	537 (10.1)
Home		
Urban	2759 (91.4)	4270 (80.7)
Rural	261 (8.6)	1023 (19.3)
Socioeconomic quintile†		
5 (highest)	250 (8.3)	705 (13.3)
4	356 (11.8)	853 (16.1)
3	517 (17.1)	1034 (19.5)
2	716 (23.7)	1067 (20.2)
1 (lowest)	1181 (39.1)	1634 (30.9)
Past year health care		
Hospital admission		
Yes	92 (3.0)	119 (2.2)
No	2928 (97.0)	5174 (97.8)
Emergency visit		
Yes	1087 (36.0)	2095 (39.6)
No	1933 (64.0)	3198 (60.4)
≥ 7 outpatient visits		
Yes	342 (11.3)	509 (9.6)
No	2678 (88.7)	4784 (90.4)
Substance use disorder diagnosis‡		
Yes	221 (7.3)	289 (5.5)
No	2799 (92.7)	5004 (94.5)
Mental health diagnosis§		
Yes	698 (23.1)	1086 (20.5)
No	2322 (76.9)	4207 (79.5)
Acute incident		
Weapon		
Hand gun	63 (2.1)	238 (4.5)
Long gun	25 (0.8)	422 (8.0)
Uncertain	2932 (97.1)	4633 (87.5)
Ambulance		
Yes	2055 (68.0)	1284 (24.3)
No	965 (32.0)	4009 (75.7)
<p>Note: OHIP = Ontario Health Insurance Plan. *Denotes self-inflicted (codes X6-, X7-, X80 - X84) or assault (codes X85 - X89, X9-, Y0-). †Based on Statistics Canada algorithm. ‡OHIP diagnostic codes 303 to 304. §OHIP diagnostic codes 290 to 316 (except 293, 294, 303, 304, 308, 310, 312).</p>		

Table 2: Acute hospital care in 8313 patients with intentional or unintentional gun injury who received emergency medical care during the study period

Characteristic	No. (%) of patients*	
	Intentional n = 3020	Unintentional n = 5293
Hospital admission		
Yes	1638 (54.2)	792 (15.0)
No	1382 (45.8)	4501 (85.0)
Surgical procedure		
Yes	1060 (35.1)	538 (10.2)
No	578 (19.1)	254 (4.8)
No admission	1382 (45.8)	4501 (85.0)
Critical care unit		
Yes	756 (25.0)	266 (5.0)
No	882 (29.2)	526 (9.9)
No admission	1382 (45.8)	4501 (85.0)
Blood transfusion†		
Yes	457 (15.1)	176 (3.3)
No	1181 (39.1)	616 (11.6)
No admission	1382 (45.8)	4501 (85.0)
Time in hospital		
No. of days, median (IQR)	6 (3–11)	5 (2–10)
No admission	–	–
Outcome‡		
Death	363 (12.0)	204 (3.9)
Alive	2657 (88.0)	5089 (96.1)

Note: IQR = interquartile range.
 *Unless stated otherwise.
 †Based on packed red cells, platelets or whole blood.
 ‡Excludes deaths at the scene and includes short-term emergency and hospital care.

10 years and found a 15% risk of hospital mortality after gun injury.⁶⁴ In contrast, the relatively lower 7% hospital mortality observed in our study may be partially attributable to a smaller number of intentional incidents with reduced injury severity. Additional studies of mortality spanning similar long-term follow-up show a comparable risk of long-term mortality despite the overall decline in hospital case fatality rates.^{65,66} Readmissions for delayed complications are also commonly reported in case series analyses of patients after gun injury, with a majority occurring after the first year postdischarge.⁶²

Systematic investigations of quality of life after gun injury are relatively sparse, typically based on small select samples, and tend to use qualitative survey responses. One study conducted by telephone interviews years after hospital discharge found that about half the patients had physical limitations and nearly two-thirds experienced psychological illnesses following gun injury.⁶⁷ The risks of long-term disability were further accentuated in patients of lower socioeconomic status.⁶⁸

These self-reported findings agree with the high prevalence of long-term disability observed in our study. To our knowledge, there are no systematic studies of long-term disability after gun injury in a Canadian population, and international literature repeatedly highlights the gaps in the scientific understanding of long-term outcomes for patients after gun injury.^{69,70}

The uncertain reasons underlying long-term disability suggest the need for a multimodal approach to trauma survivorship.⁷¹ This might include follow-up care from surgeons, psychiatrists, family physicians, physiotherapists, social workers, occupational therapists, spiritual care workers and other allied professionals.⁷² Current clinical care does not abandon patients; however, our study suggests that the prevailing efforts are not sufficient. Future clinical priorities could include additional management of pain, depression, anxiety, sleep and substance use disorder.^{73–75} Additional counselling might also be necessary for workforce participation, including a role for a navigator to negotiate between a disabled survivor and a potential employer.⁷⁶ To the best of our knowledge, no trauma centre offers such a holistic follow-up clinic for adults after gun injury.^{77,78} Our study highlights a substantial need to support survivors.⁷⁹

Future research based on additional data sources might be justified as opportunities to address these and other uncertainties.^{80,81} These studies might include broad surveys of clinical treatments and unmet health care needs over time.⁸² An alternative approach can be an evidence-based social network analysis of social dynamics among survivors over time.⁸³ Further investigations might examine specific risk factors for gun injury and predictive factors for recovery outcomes.⁸⁴ A more controversial line of study can test the subjective nature of disability and the fallible determinations of severity.⁸⁵ Additional opportunities might also include analyses of governmental and consumer databases from other countries.⁸⁶

Limitations

Our study has many limitations. The patients were mostly injured in isolated incidents and this provides little insight about gang violence or mass casualty events.^{87–89} The data were based on a large high-income region and may underestimate losses in low-resource settings. Our data may also underestimate losses owing to fallibility in identifying intentional injuries.³ The study does not account for patients who died at the scene, those who moved away (and were assumed fine) or those harmed by a culture of fear.^{70,90} Disability data may also underestimate additional losses from missed days at work, career advancement, family relationships, emotional distress, trauma recidivism and financial costs.^{91,92} The design was an observational analysis and cannot establish whether a correlation indicates possible causality (E-value = 2.80).^{93–96}

The findings may be also prone to misquotation in public debates about firearms.⁹⁷ Specifically, the observed risks of long-term disability do not directly prove or disprove calls for gun control.⁹⁸ The study does not address safe storage, background checks, licensing policies, waiting periods, safety training or other particular injury prevention strategies.^{99,100}




Figure 1: Risk of subsequent disability. Cumulative incidence plots of absolute risk of disability following injury.




Figure 2: Risk of subsequent death. Cumulative incidence plots of absolute risk of death following injury.

Table 3: Predictors of disability after gun violence*

Characteristic	Relative risk (95% CI)	
	Basic analysis†	Adjusted analysis‡
Acute injury		
Intentional injury	2.01 (1.80–2.25)	1.40 (1.24–1.60)
Age group		
Younger (Ref. ≥ 30 yr)	0.96 (0.85–1.07)	§
Sex		
Male	0.95 (0.79–1.14)	§
Home location (Ref. = rural)		
Urban	1.86 (1.54–2.25)	1.48 (1.22–1.80)
Socioeconomic quintile (Ref. = middle)		
Highest	0.81 (0.67–0.98)	0.86 (0.70–1.04)
Lowest	1.52 (1.30–1.78)	1.39 (1.19–1.63)
Incident season (Ref. = summer)		
Autumn	0.91 (0.79–1.06)	§
Winter	1.05 (0.90–1.24)	§
Spring	1.04 (0.89–1.21)	§
Incident day (Ref. = weekday)		
Weekend	1.07 (0.95–1.20)	§
Incident hour (Ref. = afternoon)		
Morning	1.09 (0.92–1.28)	1.03 (0.87–1.22)
Night	1.32 (1.17–1.50)	1.08 (0.95–1.23)
Ambulance transport		
Yes	2.21 (1.98–2.47)	1.76 (1.55–1.99)
History		
Hospital admission in past		
Yes	1.33 (0.90–1.95)	§
Emergency visit in past		
Yes	1.23 (1.10–1.38)	1.16 (1.03–1.31)
Outpatient visits in past		
Count	1.02 (1.02–1.03)	1.01 (1.00–1.01)
Mental health diagnosis in past		
Yes	1.92 (1.70–2.17)	1.67 (1.47–1.91)
Substance abuse in past		
Yes	2.43 (2.04–2.89)	1.62 (1.32–1.99)
Note: CI = confidence interval, Ref. = referent. *Estimates based on Fine and Gray model. Analyses based on all patients surviving acute gun violence. †No adjustment for baseline differences. ‡Adjusted for all measures significant in univariable analysis. §Denotes estimates not significant in univariable or multivariable model.		

The analysis provides no data on the benefits of gun ownership for those who gain security and experience no adverse incidents.¹⁰¹ The specific data on determining firearm type are difficult to interpret because of the large number of cases with missing information. The research cannot settle the unceasing

tension between safety and liberty because moral imperatives extend beyond a biomedical perspective.^{102–104}

An additional caveat is that our data do not explain mechanisms by which gun injury might lead to higher disability after intentional rather than unintentional incidents. Many

biomedical and social factors could contribute to these mechanisms including severity of injury, preexisting illnesses and background community supports.^{105–108} Another potential contributor is the fear of being labelled from identifying as a victim, which may lead to reduced self-esteem, decreased self-efficacy and depression.^{75,109–111} A further possibility may be the nature of blame, concepts of responsibility and a negative spiral of disempowerment with anxiety.^{112–114} A different explanation might be a statistical artifact from fallible ascertainment of intent leading to misclassification bias.

Conclusion

Gun injury causes considerable mortality yet many patients survive with long-term disability. The long-term risks of disability are particularly accentuated for survivors after intentional injuries.

References

1. Why own a gun? Protection is now top reason. Washington (D.C.): Pew Research Center; 2013 Mar. 12. Available: www.people-press.org/2013/03/12/why-own-a-gun-protection-is-now-top-reason/ (accessed 2019 Sept. 1).
2. Cotter A. *Firearms and violent crime in Canada, 2012*. Ottawa: Statistics Canada; 2012. Cat no 85-002-X. Available: <https://www150.statcan.gc.ca/n1/pub/85-002-x/2014001/article/11925-eng.htm> (accessed 2019 Feb. 12).
3. Yancher NL, Beno S; Canadian Association of Emergency Physicians and the Trauma Association of Canada. Can we do better?: A Canadian perspective on firearm injury prevention. *Ann Surg* 2018;267:1009–10.
4. Nand D, Naghavi M, Marczak LB, et al. Global mortality from firearms, 1990–2016. *JAMA* 2018;320:792–814.
5. WHO Mortality Data Base. Geneva: World Health Organization Department of Health Statistics and Information Systems; updated 2018 May. Available: http://apps.who.int/healthinfo/statistics/mortality/causeofdeath_query/ (accessed 2019 Feb. 12).
6. Corso PS, Mercy JA, Simon TR, et al. Medical costs and productivity losses due to interpersonal and self-directed violence in the United States. *Am J Prev Med* 2007;32:474–82.
7. Raza S, Redelmeier DA. Gunshot to the head. *Am J Med* 2018;131:e7–8.
8. Ajmal S, Enam SA, Shamim MS. Neurogenic claudication and radiculopathy as delayed presentations of retained spinal bullet. *Spine J* 2009;9:e5–8.
9. Wiseman T, Foster K, Curtis K. Mental health following traumatic physical injury: an integrative literature review. *Injury* 2013;44:1383–90.
10. Disaster Technical Assistance Center Supplemental Research Bulletin: Mass violence and behavioral health. Rockville (MD): Substance Abuse and Mental Health Services Administration; 2017:1–18. Available: <https://www.samhsa.gov/sites/default/files/dtac/srb-mass-violence-behavioral-health.pdf> (accessed 2019 July 26).
11. Smith RN, Seamon MJ, Kumar V, et al. Lasting impression of violence: retained bullets and depressive symptoms. *Injury* 2018;49:135–40.
12. Lee J. Wounded: life after the shooting. *Ann Am Acad Pol Soc Sci* 2012;642:244–57.
13. Eligon J. One bullet can kill, but sometimes 20 don't, survivors show. *The New York Times* 2008 Apr. 3. Available: <https://www.nytimes.com/2008/04/03/nyregion/03shot.html> (accessed 2019 July 26).
14. Ringburg AN, Polinder S, Van Ierland MCP, et al. Prevalence and prognostic factors of disability after major trauma. *J Trauma* 2011;70:916–22.
15. Laughlin MD, Belmont PJ, Lanier PJ, et al. Occupational outcomes following combat-related gunshot injury. *Int J Surg* 2017;48:286–90.
16. Stevenson T, Carr DJ, Penn-Barwell JG, et al. The burden of gunshot wounding of UK military personnel in Iraq and Afghanistan from 2003–14. *Injury* 2018;49:1064–9.
17. Redelmeier DA, Blair PJ. Survivors of motor vehicle trauma — an analysis of seat belt use and health care utilization. *J Gen Intern Med* 1993;8:199–203.
18. Mills BM, Nurius PS, Matsueda RL, et al. Prior arrest, substance use, mental disorder, and intent-specific firearm injury. *Am J Prev Med* 2018;55:298–307.
19. Rattan R, Parreco J, Namias N, et al. Hidden costs of hospitalization after firearm injury: national analysis of different hospital readmission. *Ann Surg* 2018;267:810–5.
20. *Canada at a glance 2010 — population*. Ottawa: Statistics Canada; 2010. Available: <https://www150.statcan.gc.ca/n1/pub/12-581-x/2010000/pop-eng.htm> (accessed 2019 Mar. 17).
21. About Ontario. Toronto: Government of Ontario; 2019. Available: <https://www.ontario.ca/page/about-ontario> (accessed 2019 Mar. 17).
22. van Walraven CV, Paterson JM, Kapral M, et al. Appropriateness of primary total hip and knee replacements in regions of Ontario with high and low utilization rates. *CMAJ* 1996;155:697–706.
23. Iron K, Manuel DG. Quality assessment of administrative data (QuAAD): an opportunity for enhancing Ontario's health data. Toronto: ICES; 2007.
24. Iron K, Zagorski B, Sykora K. Living and dying in Ontario: an opportunity for improved health information — ICES investigative report. Toronto: ICES; 2008. Available: <https://www.ices.on.ca/Publications/Atlases-and-Reports/2008/Living-and-dying-in-Ontario> (accessed 2019 July 26).
25. Chiu M, Lebenbaum M, Lam K, et al. Describing the linkages of the immigration, refugees and citizenship Canada permanent resident data and vital statistics death registry to Ontario's administrative health database. *BMC Med Inform Decis Mak* 2016;16:135.
26. Data dictionary. Toronto: ICES; 2019. Available: <https://datadictionary.ices.on.ca/Applications/DataDictionary/Default.aspx> (accessed 2020 Feb. 11).
27. Working with ICES data. Toronto: ICES. Available: www.ices.on.ca/Data-and-Privacy/ICES-data/Working-with-ICES-Data (accessed 2020 Feb. 11).
28. Chan B, Schull MJ, Schultz SE. Atlas report: emergency department services in Ontario. Toronto: ICES; 2001. Available: <https://www.ices.on.ca/flip-publication/emergency-department-services-in-ontario/files/assets/basic-html/index.html#54> (accessed 2019 July 26).
29. NACRS emergency department visits and length of stay, 2017–2018 [National Ambulatory Care Reporting System]. Ottawa: Canadian Institute for Health Information; 2019. Available: <https://www.cihi.ca/en/quick-stats> (accessed 2019 Mar. 17).
30. Kondro W. Gunshot wound reporting mandatory in Ontario. *CMAJ* 2005;173:242–242-a.
31. Ministry of Community and Social Services. Ontario Ministry of Children, Community and Social Services. Toronto; 2019. Available: <https://www.mcscs.gov.on.ca/en/mcscs/> (accessed 2019 Mar. 17).
32. Morris S, Fawcett G, Brisebois L, et al. *A demographic, employment and income profile of Canadians with disabilities aged 15 years and over, 2017*. Ottawa: Statistics Canada; 2018. Cat no 89-654-X2018002. Available: <https://www150.statcan.gc.ca/n1/pub/89-654-x/89-654-x2018002-eng.pdf> (accessed 2019 July 26).
33. Data quality documentation, National Ambulatory Care Reporting System: current-year information, 2017–2018. Ottawa: Canadian Institute for Health Information; 2018. Available: <https://www.cihi.ca/sites/default/files/document/current-year-information-nacrs-2017-2018-en-web.pdf> (accessed 2019 July 26).
34. Data quality study of Ontario emergency department visits for 2004–2005 — executive summary. Ottawa: Canadian Institute for Health Information; 2007:1–22. Available: https://secure.cihi.ca/free_products/vol1_nacrs_executive_summary_nov2_2007.pdf (accessed 2019 July 26).
35. LeMier M, Cummings P, West TA. Accuracy of external cause of injury codes reported in Washington State hospital discharge records. *Inj Prev* 2001;7:334–8.
36. Juurlink D, Preyra C, Croxford R, et al. Canadian institute for health information discharge abstract database: a validation study. Toronto: ICES; 2006.
37. Macpherson AK, Schull MJ. Penetrating trauma in Ontario emergency departments: a population-based study. *CJEM* 2007;9:16–20.
38. Furlan JC, Fehlings MG. The National Trauma Registry as a Canadian spine trauma database: a validation study using an institutional clinical database. *Neuroepidemiology* 2011;37:96–101.
39. Altoijry A, Al-Omran M, Lindsay TF, et al. Validity of vascular trauma codes at major trauma centres. *Can J Surg* 2013;56:405–8.
40. Hulley SB, Cummings SR. *Designing clinical research*. Baltimore: Williams and Wilkins; 1988.
41. *Postal code conversion file (PCCF), reference guide*. Ottawa: Statistics Canada; 2017. Cat no 92-154-G. Available: www.canadapost.ca/cpc/assets/cpc/uploads/files/marketing/2017-postal-code-conversion-file-reference-guide-en.pdf (accessed 2019 Feb. 11).
42. *Income and earnings reference guide, 2006 census*. Ottawa: Statistics Canada; 2006. Cat no 97-563-GWE2006003. Available: <http://www12.statcan.gc.ca/census-recensement/2006/ref/rp-guides/income-revenu-eng.cfm> (accessed 2019 July 26).
43. Parachute. Ontario injury data report 2018. Toronto: Parachute; 2018.
44. Becker R, Silvi J, Ma Fat D, et al. A method for deriving leading causes of death. *Bull World Health Organ* 2006;84:297–304.
45. Ardal S. Health analyst's toolkit. Toronto: Ontario Ministry of Health and Long-Term Care; 2012. Available: http://www.health.gov.on.ca/english/providers/pub/healthanalytics/health_toolkit/health_toolkit.pdf (accessed 2019 July 26).
46. Discharge abstract database. Ottawa: Canadian Institute for Health Information; 2019. Available: <https://www.cihi.ca/en/discharge-abstract-database-metadata> (accessed 2019 July 26).
47. Drucker AM, Thiruchelvan D, Redelmeier DA. Eczema and subsequent suicide: a matched case-control study. *BMJ Open* 2018;8:e023776.
48. Newgard CD, Sanchez BJ, Bulger EM, et al. A geospatial analysis of severe firearm injuries compared to other injury mechanisms: event characteristics, location, timing, and outcomes. *Acad Emerg Med* 2016;23:554–65.
49. Gani F, Sakran JV, Canner JK. Emergency department visits for firearm-related injuries in the United States, 2006–14. *Health Aff (Millwood)* 2017;36:1729–38.
50. Sauaia A, Gonzalez E, Moore HB, et al. Fatality and severity of firearm injuries in a denver trauma center, 2000–2013. *JAMA* 2016;315:2465–7.

51. Saunders NR, Lee H, Macpherson A, et al. Risk of firearm injuries among children and youth of immigrant families. *CMAJ* 2017;189:E452-8.
52. Kapral MK, Fang J, Chan C, et al. Neighborhood income and stroke care and outcomes. *Neurology* 2012;79:1200-7.
53. Riddell CA, Harper S, Cerda M, et al. Comparison of rates of firearm and non-firearm homicide and suicide in black and white non-Hispanic men, by U.S. State. *Ann Intern Med* 2018;168:712-20.
54. Ontario Health Insurance Plan. Toronto: Ontario Ministry of Health and Long-Term Care; 2016. Available: <http://www.health.gov.on.ca/en/public/programs/ohip/> (accessed 2016 Oct. 12).
55. Ontario Health Insurance Plan: OHIP schedule of benefits and fees. Toronto: Government of Ontario Ministry of Health and Long-Term Care; 2018. Available: <http://www.health.gov.on.ca/en/pro/programs/ohip/sob/> (accessed 2019 Mar. 17).
56. Reece SCM, Thiruchelvam D, Redelmeier DA. Medical emergencies in farmers. *J Rural Health* 2019;35:429-35.
57. Austin PC, Lee DS, Fine JP. Introduction to the analysis of survival data in the presence of competing risks. *Circulation* 2016;133:601-9.
58. Fine JP, Gray RJ. A proportional hazards model for the subdistribution of a competing risk. *J Am Stat Assoc* 1999;94:496-509.
59. Putter H, Fiocco M, Geskus RB. Tutorial in biostatistics: competing risks and multi-state models. *Stat Med* 2007;26:2389-430.
60. Lau B, Cole SR, Gange SJ. Competing risk regression models for epidemiologic data. *Am J Epidemiol* 2009;170:244-56.
61. Austin PC, Fine JP. Practical recommendations for reporting Fine-Gray model analyses for competing risk data. *Stat Med* 2017;36:4391-400.
62. Wintemute GJ, Wright MA. Initial and subsequent hospital costs of firearm injuries. *J Trauma* 1992;33:556-60.
63. Gabbe BJ, Simpson PM, Harrison JE, et al. Return to work and functional outcomes after major trauma who recovers, when, and how well? *Ann Surg* 2016;263:623-32.
64. Haider AH, Young JH, Kisat M, et al. Association between intentional injury and long-term survival after trauma. *Ann Surg* 2014;259:985-92.
65. Davidson GH, Hamlat CA, Rivara FP, et al. Long-term survival of adult trauma patients. *JAMA* 2011;305:1001-7.
66. Fahimi J, Larimer E, Hamud-Ahmed W, et al. Long-term mortality of patients surviving firearm violence. *Inj Prev* 2016;22:129-34.
67. Vella MA, Warshauer A, Tortorello G, et al. Long-term functional, psychological, emotional, and social outcomes in survivors of firearm injuries. *JAMA Surg* 2019;155:51-9.
68. Herrera-Escobar JP, Seshadri AJ, Rivero R, et al. Lower education and income predict worse long-term outcomes after injury. *J Trauma Acute Care Surg* 2019;87:104-10.
69. Bulger EM, Kuhls DA, Campbell BT, et al. Proceedings from the medical summit on firearm injury prevention: a public health approach to reduce death and disability in the US. *J Am Coll Surg* 2019;229:415-30.e12.
70. Rowhani-Rahbar A, Zatzick DF, Rivara FP. Long-lasting consequences of gun violence and mass shootings. *JAMA* 2019;321:1765-6.
71. Hobfoll SE, Watson P, Bell CC, et al. Five essential elements of immediate and mid-term mass trauma intervention: empirical evidence. *Psychiatry* 2007;70:283-315.
72. Walter E, Curtis K. The role and impact of the specialist trauma nurse: an integrative review. *J Trauma Nurs* 2015;22:153-69.
73. Lee J. The pill hustle: risky pain management for a gunshot victim. *Soc Sci Med* 2013;99:162-8.
74. Wade D, Crompton D, Howard A, et al. Skills for psychological recovery: evaluation of a post-disaster mental health training program. *Disaster Health* 2015;2:138-45.
75. Patton D, Sodhi A, Affinati S, et al. Post-discharge needs of victims of gun violence in Chicago: a qualitative study. *J Interpers Violence* 2019;34:135-55.
76. Shaw W, Hong QN, Pransky G, et al. A literature review describing the role of return-to-work coordinators in trial programs and interventions designed to prevent workplace disability. *J Occup Rehabil* 2008;18:2-15.
77. Khan F, Amatya B, Hoffman K. Systematic review of multidisciplinary rehabilitation in patients with multiple trauma. *Br J Surg* 2012;99:88-96.
78. Taichman DB, Bauchner H, Drazen JM, et al. Firearm-related injury and death: a U.S. Health care crisis in need of health care professionals. *PLoS Med* 2017;14:e1002430.
79. Stanbrook MB. Gun control: a health issue for which physicians rightfully advocate. *CMAJ* 2019;191:E434-5.
80. Bingenheimer JB, Brennan RT, Earls FJ. Firearm violence exposure and serious violent behavior. *Science* 2005;308:1323-6.
81. Spitzer SA, Vail D, Tennakoon L, et al. Readmission risk and costs of firearm injuries in the United States, 2010-2015. *PLoS One* 2019;14:e0209896.
82. Casey R. Disability and unmet health care needs in Canada: a longitudinal analysis. *Disabil Health J* 2015;8:173-81.
83. Green B, Horel T, Papachristos AV. Modeling contagion through social networks to explain and predict gunshot violence in Chicago, 2006 to 2014. *JAMA Intern Med* 2017;177:326-33.
84. Branas CC, Han S, Wiebe DJ. Alcohol use and firearm violence. *Epidemiol Rev* 2016;38:32-45.
85. Steenstra IA, Busse JW, Tulosso D, et al. Predicting time on prolonged benefits for injured workers with acute back pain. *J Occup Rehabil* 2015;25:267-78.
86. National Electronic Injury Surveillance System (NEISS). Bethesda (MD): U.S. Consumer Products Safety Commission; 2014. Available: http://www.cpsc.gov/Global/Neiss_prod/completemanual%20.pdf (accessed 2019 July 26).
87. Studdert DM, Zhang Y, Rodden JA, et al. Handgun acquisitions in California after two mass shootings. *Ann Intern Med* 2017;166:698-706.
88. Wintemute GJ. How to stop mass shootings. *N Engl J Med* 2018;379:1193-6.
89. Flegler E. Mass shootings and the numbing of America. *JAMA Intern Med* 2019;179:610-1.
90. Larsen DA, Lane S, Jennings-Bey T, et al. Spatio-temporal patterns of gun violence in Syracuse, New York 2009-2015. *PLoS One* 2017;12:e0173001.
91. Diez C, Kurland RP, Rothman EF, et al. State intimate partner violence-related firearm laws and intimate partner homicide rates in the United States, 1991 to 2015. *Ann Intern Med* 2017;167:536-43.
92. The cost of injury in Canada. Toronto: Parachute; 2015.
93. VanderWeele TJ, Ding P. Sensitivity analysis in observational research: introducing the E-value. *Ann Intern Med* 2017;167:268-74.
94. VanderWeele TJ, Mathur MB, Ding P. Correcting misinterpretations of the E-value. *Ann Intern Med* 2019;170:131-2.
95. Haneuse S, VanderWeele TJ, Arterburn D. Using the E-value to assess the potential effect of unmeasured confounding in observational studies. *JAMA* 2019;321:602-3.
96. Trinquart L, Erlinger AL, Petersen JM, et al. Applying the E-value to assess the robustness of epidemiologic fields of inquiry to unmeasured confounding. *Am J Epidemiol* 2019;188:1174-80.
97. Maa J, Darzi A. Firearm injuries and violence prevention — the potential power of a surgeon general's report. *N Engl J Med* 2018;379:408-10.
98. Bauchner H, Rivara FP, Bonow RO, et al. Death by gun violence — a public health crisis. *JAMA* 2017;318:1763-4.
99. Matthey EC, Galin J, Rudolph KE, et al. In-state and interstate associations between gun shows and firearm deaths and injuries. *Ann Intern Med* 2017;167:837-44.
100. Institute of Medicine, National Research Council. Priorities for research to reduce the threat of firearm-related violence. Washington (D.C.): The National Academies Press; 2013. Available: <http://www.nap.edu/read/18319> (accessed 2019 Apr. 15).
101. Campion EW, Morrissey S, Malina D, et al. After the mass shooting in Las Vegas — finding common ground on gun control. *N Engl J Med* 2017;377:1679-80.
102. Butkus R, Weissman A. Internists' attitudes toward prevention of firearm injury. *Ann Intern Med* 2014;160:821-7.
103. Malina D, Morrissey S, Campion EW, et al. Rooting out gun violence. *N Engl J Med* 2016;374:175-6.
104. Cunningham RM, Zimmerman MA, Carter PM. Money, politics and firearm safety. *JAMA NetW Open* 2019;2:e187823.
105. van Delft-Schreurs CCHM, Van Bergen JJM, De Jongh MAC, et al. Quality of life in severely injured patients depends on psychosocial factors rather than on severity or type of injury. *Injury* 2014;45:320-6.
106. Garber BG, Rusu C, Zamorski MA, et al. Occupational outcomes following mild traumatic brain injury in Canadian military personnel deployed in support of the mission in Afghanistan: a retrospective cohort study. *BMJ Open* 2016;6:e010780.
107. Goldstick JE, Carter PM, Walton MA, et al. Development of the SaFETy score: a clinical screening tool for predicting future firearm violence risk. *Ann Intern Med* 2017;166:707-14.
108. Rousseau C, Hassan G, Frounfelker R. The role of physicians in the violence epidemic. *CMAJ* 2019;191:E644.
109. Aronson E. The theory of cognitive dissonance: a current perspective. *Adv Exp Soc Psychol* 1969;4:1-34.
110. Martz E. Death anxiety as a predictor of posttraumatic stress levels among individuals with spinal cord injuries. *Death Stud* 2004;28:1-17.
111. Gollwitzer M, Süssenbach P, Hannuschke M. Victimization experiences and the stabilization of victim sensitivity. *Front Psychol* 2015;6:439.
112. Shaver KG. *The attribution of blame: causality, responsibility, and blameworthiness*. New York: Springer; 1985.
113. Liebschutz J, Schwartz S, Hoyte J, et al. A chasm between injury and care: Experiences of black male victims of violence. *J Trauma* 2010;69:1372-8.
114. Callebaut L, Molyneux P, Alexander T. The relationship between self-blame for the onset of a chronic physical health condition and emotional distress: a systematic literature review. *Clin Psychol Psychother* 2017;24:965-86.

Affiliations: Department of Medicine (Raza, Redelmeier), University of Toronto; Evaluative Clinical Sciences (Raza, Thiruchelvam, Redelmeier), Sunnybrook Research Institute; ICES in Ontario (Thiruchelvam); Institute of Health Policy, Management and Evaluation (Redelmeier), Toronto, Ont.

Contributors: Donald Redelmeier wrote the first draft. Sheharyar Raza and Donald Redelmeier contributed to study design and background scholarship. All authors contributed to manuscript preparation, data analysis, results interpretation, critical revisions and the final decision to

submit. All authors provided final approval of the version to be published. All authors agreed to be accountable for all aspects of the work.

Funding: This project was supported by a Canada Research Chair in Medical Decision Sciences, the Canadian Institutes of Health Research, and the PSI Foundation of Ontario. The funding organizations had no role in the design and conduct of the study; collection, management, analysis or interpretation of the data; and preparation, review or approval of the manuscript.

Acknowledgments: The authors thank the following for their helpful comments: Leonard Evans, Fizza Manzoor, Robert Redelmeier, Filippo Raso, Lee Ross, Charles Tator and Albert Wu.

Data sharing: The data set from this study is held securely in coded form at ICES. While data sharing agreements prohibit ICES from making the data set publicly available, access may be granted to those who meet pre-specified criteria for confidential access, available at www.ices.on.ca/DAS. The full data set creation plan and underlying analytic code are available

from the authors upon request, understanding that the computer programs may rely upon coding templates or macros that are unique to ICES and are therefore either inaccessible or may require modification. Patient privacy laws prohibit making individual-level data publicly available. Aggregate data are shown in the paper and Appendix 1. Researchers interested in replicating or extending the work can seek access to individual-level data through ICES (www.ices.on.ca).

Supplemental information: For reviewer comments and the original submission of this manuscript, please see www.cmajopen.ca/content/8/3/E469/suppl/DC1.

Disclaimer: This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health and Long-Term Care (MOHLTC). The opinions, results and conclusions reported in this article are those of the authors and are independent from the funding sources. No endorsement by ICES or the Ontario MOHLTC is intended or should be inferred.